

SCHIZOPHRENIA-LIKE RELATIVE NUTRITIONAL DEFICIENCY



SINCE 1828

differential diagnosis noun

Definition of *differential diagnosis*
: the distinguishing of a disease or condition from others presenting with similar signs and symptoms

RELATIVE NUTRITIONAL DEFICIENCY (RND)

A relative nutritional deficiency occurs when an optimal diet does not meet the needs of the system.

PROVISIONAL DIAGNOSIS: SCHIZOPHRENIA

After making the provisional diagnosis of schizophrenia, formulation of a differential diagnosis is required followed by a medical work up to determine the working diagnosis.

DIFFERENTIAL DIAGNOSIS

SCHIZOPHRENIA (PARTIAL ILLUSTRATIVE):

1. Schizophrenia
2. **Schizophrenia-like dopamine related relative nutritional deficiency symptoms™**
3. Brief psychotic disorder
4. Delirium
5. Delusional disorder
6. Medical illnesses
7. Medication-induced disorder
8. Mood disorders with psychotic features
9. Pervasive developmental disorder
10. Psychotic disorder NOS
11. Schizophreniform disorder
12. Schizotypal personality disorder
13. Substance abuse

Management of

SCHIZOPHRENIA-LIKE DOPAMINE RELATED RELATIVE NUTRITIONAL DEFICIENCY SYMPTOMS™ *

If systemic dopamine (catecholamine) concentrations are low, inadequate, depleted, or suboptimal™ on an optimal diet, a relative nutritional deficiency of their precursors or cofactors always exists. When schizophrenia-like RND symptoms are present, consider a dopamine challenge,™ of dopamine precursors along with properly balanced serotonin precursors, thiols and cofactors.

Baseline urinary and serum dopamine assays are not reproducible, therefore, are of no assistance in establishing the existence of dopamine concentrations that are not high enough or nutrient starting point for addressing relative nutritional deficiency related symptoms.

Mastery of side effect management should be in place before starting the dopamine precursor protocol.

THE UPPER DOSING RANGE

Under the dopamine related RND protocol™ standardized 40% Mucuna Pruriens is increased by 2.4 grams per week. If the development of side effects does not increase the time to stabilization, some patients may require ten to fifteen weekly Mucuna increases (24 gr. to 36 gr. per day or more) before finding the optimal individualized dosing associated with schizophrenia-like dopamine related RND symptom relief.™

* Covered by other trademarks: RND™, RND symptoms™, relative nutritional deficiency symptoms™, relative nutritional deficiency™, dopamine relative nutritional deficiency™, Schizophrenia-like dopamine relative nutritional deficiency™, dopamine RND™, like RND™, like relative nutritional deficiency™, like dopamine™,



INFO@NEUROSupport.COM



1150 88TH AVE W., DULUTH, MN



+1-218-626-2220